

Irreducible Modules for Classical and Alternating Groups

William J. Husen

metadata, citation and similar papers at core.ac.uk

Communicated by Jan Saxl

Received October 6, 1999

1. INTRODUCTION

In this article we consider the following question: Given groups $H < G$ and an absolutely irreducible $\mathbb{F}G$ -module M , when does $M \downarrow_H$ remain irreducible? This question arises naturally in the study of the maximal subgroups of the classical groups. We assume that $H \cong A_n$ or $2.A_n$, $n \geq 15$, and that G is a simply connected classical algebraic group over \mathbb{F} , where \mathbb{F} is an algebraically closed field of characteristic $p \geq 0$. In particular, $G = SL(V)$, $Sp(V)$, or $Spin(V)$, where V is a vector space over \mathbb{F} . For convenience, we assume that $\dim(V)$ is even whenever $G = Spin(V)$ and $p = 2$. In this context we prove the following theorem:

THEOREM 1.1. *Suppose that $M = M(\lambda)$ is a tensor indecomposable $\mathbb{F}G$ -module such that $M \downarrow_H$ is absolutely irreducible and $\dim(M) > \dim(V)$.*

1. *If $G = SL(V)$ then either*
 - (a) $\dim(V) \leq n^3$ or
 - (b) λ or $\rho(\lambda) = 3\lambda_1, 2\lambda_1 + \lambda_2, 2\lambda_1 + \lambda_\ell, \lambda_2 + \lambda_\ell$, or λ_3 and $\dim(V) \leq n^4$.
2. *If $G = Sp(V)$ or $Spin(V)$ then either*
 - (a) $\dim(V) \leq 2n^4$ or
 - (b) $\lambda = 3\lambda_1, 2\lambda_1 + \lambda_2$, or λ_3 and $\dim(V) \leq 2n^6$.

Furthermore, if $n \geq 74$ and $p \neq 2$, then $G \neq Sp(V)$.



This result strengthens the main result of [3]. Throughout we will assume that λ is the highest weight of M as an $\mathbb{F}G$ -module and that M satisfies the hypothesis of Theorem 1.1. As M is tensor indecomposable, we may replace M with a Galois conjugate and assume that λ is a p -restricted dominant weight when $p > 0$. Then we proceed as follows: Using a result of Premet, suitably extended, we find a vector of weight μ in M which has a small stabilizer W_μ in the Weyl group W of G . Then $[W : W_\mu] \leq \dim(M)$. Conversely, we produce a large subgroup \tilde{C} of H such that $M \downarrow_{\tilde{C}}$ contains a small submodule M_0 . Using Frobenius reciprocity, we get $\dim(M) \leq \dim(M_0)[H : \tilde{C}]$. The result follows from comparing these bounds.

Our result is analogous to that achieved in [5] for H a classical group. In similar notation, they find an upper bound for $\dim(M)$ in terms of $\dim(V)$. This is in contrast to our result, where we find an upper bound for $\dim(V)$ in terms of n . This reflects the fact that there is an infinite family of examples where $\dim(V) < n$ and $M \downarrow_H$ is irreducible. In particular, if V is the minimal $n-1$ or $n-2$ dimensional module for H over \mathbb{F} , $G = Spin(V)$ or $SL(V)$, and $M = \wedge^k V$, $k < \frac{n}{2}$, then $M \downarrow_H$ is absolutely irreducible.

2. NOTATION AND PRELIMINARY RESULTS

We use the following notation throughout this article:

ℓ is the Lie rank of G and $e = \lfloor \log_n(\ell) \rfloor$.

$\bar{e} = e$ or $\frac{3e}{2}$ if $e \leq 3$ or $e > 3$, respectively.

T is a fixed maximal torus of G .

If $G = Spin(V)$ or $Sp(V)$, we write $t = (t_1, \dots, t_\ell)$ for an arbitrary $t \in T$.

If $G = SL(V)$, we write $t = (t_1, \dots, t_{\ell+1})$ for $t \in T$, considering T as a subgroup of $GL(V)$.

$\Pi = \{\alpha_1, \dots, \alpha_\ell\}$ are the simple roots of G with respect to T , ordered as in [2].

Φ is the root system generated by Π and Φ^+ is the set of positive roots.

$\{\lambda_1, \dots, \lambda_\ell\}$ are the fundamental dominant weights of G ordered according to the ordering of Π . For convenience, we define $\lambda_0 = \lambda_{\ell+1} = 0$.

If $G = SL(V)$ then ρ is the permutation of Π and $\{\lambda_1, \dots, \lambda_\ell\}$ induced by the non-trivial symmetry of the Dynkin diagram of G .

M_μ is the μ -weight space of M .

Λ is the weight lattice of G and Λ^+ is the set of dominant weights in Λ .

$\lambda = \sum_{i=1}^{\ell} a_i \lambda_i$ is the highest weight of M .

$k_s = \sum_{i < \ell/2} i a_i$, $k_d = \sum_{i \geq \ell/2} (\ell - i + 1) a_i$, and $k_d^* = \ell + 1 - k_d$.

W is the Weyl group of G and W_μ is the stabilizer of $\mu \in \Lambda$ in W .

ϵ_i is the element of $\text{Hom}(T, F^*)$ such that $\epsilon_i(t) = t_i$.

It will be convenient to know α_i and λ_i in terms of ϵ_i :

$G = SL(V)$:

$$\alpha_i = \epsilon_i - \epsilon_{i+1}, \quad \lambda_i = \sum_{j=1}^i \epsilon_j.$$

$G = Sp(V)$:

$$\alpha_i = \begin{cases} \epsilon_i - \epsilon_{i+1}, & i < \ell, \\ 2\epsilon_\ell, & i = \ell, \end{cases} \quad \lambda_i = \sum_{j=1}^i \epsilon_j.$$

$G = Spin(V)$, $\dim(V)$ odd:

$$\alpha_i = \begin{cases} \epsilon_i - \epsilon_{i+1}, & i < \ell, \\ \epsilon_\ell, & i = \ell, \end{cases} \quad \lambda_i = \begin{cases} \sum_{j=1}^i \epsilon_j, & i < \ell, \\ \frac{1}{2} \sum_{j=1}^{\ell} \epsilon_j, & i = \ell. \end{cases}$$

$G = Spin(V)$, $\dim(V)$ even:

$$\alpha_i = \begin{cases} \epsilon_i - \epsilon_{i+1}, & i < \ell, \\ \epsilon_{\ell-1} + \epsilon_\ell, & i = \ell, \end{cases} \quad \lambda_i = \begin{cases} \sum_{j=1}^i \epsilon_j, & i < \ell - 1, \\ \frac{1}{2} \sum_{j=1}^{\ell} \epsilon_j, & i = \ell - 1, \\ \frac{1}{2} \left(\sum_{j=1}^{\ell-1} \epsilon_j - \epsilon_\ell \right), & i = \ell. \end{cases}$$

Note that, by considering $T < GL(V)$, we have a simpler characterization of λ_i in terms of the coordinate functions ϵ_i when $G = SL(V)$.

For $\mu \in \Lambda^+$, write $\mu = \sum_{i=1}^{\ell} b_i \lambda_i$ and define $m_s(\mu) = \max\{0, i \mid i < \frac{\ell}{2} \text{ and } b_i \neq 0\}$, $m_d(\mu) = \max\{0, \ell - i + 1 \mid i \geq \frac{\ell}{2} \text{ and } b_i \neq 0\}$, and $m_d^*(\mu) = \ell + 1 - m_d(\mu)$.

If U is an \mathbb{F} -vector space, we write $U^{\otimes i} = \bigotimes_{j=1}^i U$ and define $U^{\otimes 0} = \mathbb{F}$.

$\{v_i\}$ is a basis for V such that $t \in T$ acts on this basis as $\text{diag}(t_1, \dots, t_{\ell+1})$ if $G = SL(V)$, as $\text{diag}(t_1, \dots, t_\ell, t_\ell^{-1}, \dots, t_1^{-1})$ if $G = Sp(V)$ or $Spin(V)$ with $\dim(V)$ even, and as $\text{diag}(t_1, \dots, t_\ell, 1, t_\ell^{-1}, \dots, t_1^{-1})$ if $G = Spin(V)$ with $\dim(V)$ odd.

If $G = SL(V)$, then $\{v_i^*\}$ is a basis for V^* such that $tv_i^* = t_i^{-1}v_i^*$. Note that $v_{\ell+1}^*$ is a vector of weight λ_ℓ .

Our initial result concerns binomial coefficients:

LEMMA 2.1. *Suppose that i_1 , i_2 , and j are integers.*

1. *If $1 \leq i_1 \leq j$ then*

$$\binom{j}{i_1} \geq \left(\frac{j - i_1 + 1}{i_1} \right)^{i_1}.$$

2. *If $e \geq 3$, $j \geq n^e$, and $\frac{n}{e} \leq i_1 \leq \frac{j}{2}$ then $\binom{j}{i_1} > \dim(M)$.*

3. *If $0 \leq i_1 \leq i_2 \leq j$ then*

$$\frac{j!}{i_1!i_2!(j - i_1 - i_2)!} \geq \binom{j}{i_1 + i_2}.$$

Proof. Part (1) follows by writing

$$\binom{j}{i_1} = \prod_{m=1}^{i_1} \frac{j - m + 1}{i_1 - m + 1} \geq \prod_{m=1}^{i_1} \frac{j - i_1 + 1}{i_1} = \left(\frac{j - i_1 + 1}{i_1} \right)^{i_1}.$$

For (2), $\frac{n}{e} \leq i_1 \leq \frac{j}{2}$ and $j \geq n^e$ implies that

$$\binom{j}{i_1} \geq \binom{j}{n/e} \geq \left(\frac{n^e}{n/e} \right).$$

Applying (1) to the last expression gives

$$\binom{j}{i_1} \geq \left(\frac{n^e - n/e + 1}{n/e} \right)^{n/e}.$$

As $e \geq 3$, we have

$$n^e - \frac{n}{e} + 1 > \frac{n^e}{e} \quad \text{and} \quad \frac{e - 1}{e} \geq \frac{1}{2}.$$

This gives

$$\binom{j}{i_1} > \left(\frac{n^e/e}{n/e} \right)^{n/e} \geq n^{n/2}.$$

Now, M is an irreducible $\mathbb{F}H$ -module, so $\dim(M)$ is bounded above by $\chi(1)$ for some $\chi \in \text{Irr}(H)$. Using the orthogonality of such characters, $\chi(1) < \sqrt{|H|} \leq \sqrt{n!} < n^{n/2}$, hence $\dim(M) < n^{n/2}$. This proves (2). Part (3) follows by expanding

$$\binom{j}{i_1 + i_2} = \frac{j!}{(i_1 + i_2)!(j - i_1 - i_2)!}$$

and comparing denominators. In particular, $(i_1 + i_2)! \geq i_1!i_2!$, hence the result. ■

LEMMA 2.2. *Suppose that $e \geq 3$ and $\mu \in \Lambda^+$. If M possesses a vector of weight μ , then $m_s(\mu) + m_d(\mu) < \frac{n}{e}$. Furthermore, $m_d(\mu) = 0$ if $G = \text{Sp}(V)$ or $\text{Spin}(V)$.*

Proof. Write $\mu = \sum_{i=1}^{\ell} b_i \lambda_i$. If $b_i > 0$ then $W_{\mu} \leq W_{\lambda_i}$, which implies that $\dim(M) \geq [W : W_{\mu}] \geq [W : W_{\lambda_i}]$. Let $G = Sp(V)$ or $Spin(V)$ so that $[W : W_{\lambda_i}] = 2^i \binom{\ell}{i}$, $2^{\ell-3}(\ell^2 - \ell)$, or $2^{\ell-1}$, the latter two holding if and only if $G = Spin(V)$ with $\dim(V)$ even and $i = \ell - 2$ or $i > \ell - 2$, respectively. If $0 \neq m_d(\mu) \leq \ell - 3$ then $b_{m_d^*(\mu)} > 0$ and $\dim(M) \geq 2^{m_d^*(\mu)} \binom{\ell}{m_d^*(\mu)} \geq 2^{m_d^*(\mu)} \geq 2^{\ell/2} \geq 2^{n^3/2} > \dim(M)$, which is a contradiction. Similar calculations handle the case $m_d(\mu) \geq \ell - 2$. Therefore $m_d(\mu) = 0$ and $m_s \neq 0$, so that $\dim(M) \geq 2^{m_s(\mu)} \binom{\ell}{m_s(\mu)} \geq \binom{\ell}{m_s(\mu)}$. Since $m_s(\mu) \leq \frac{\ell}{2}$, part (2) of Lemma 2.1 implies that $\binom{\ell}{m_s(\mu)} > \dim(M)$ if $m_s(\mu) \geq \frac{n}{e}$, hence $m_s(\mu) < \frac{n}{e}$.

Now suppose that $G = SL(V)$, so that $[W : W_{\lambda_i}] = \binom{\ell+1}{i}$. Since $m_d^*(\mu) = \ell + 1 - m_d(\mu)$, we have $\binom{\ell+1}{m_d^*(\mu)} = \binom{\ell+1}{m_d(\mu)}$ and $m_d(\mu) \leq \frac{\ell}{2}$. Arguing as before, $\binom{\ell+1}{m_d(\mu)} > \dim(M)$ if $m_d(\mu) \geq \frac{n}{e}$. Thus $m_d(\mu) < \frac{n}{e}$. Similarly, $m_s(\mu) < \frac{n}{e}$. Thus, as $e \geq 3$, $m_s(\mu) + m_d(\mu) < \frac{\ell}{2}$. Now $W_{\mu} < W_{\lambda_{m_s(\mu)}} \cap W_{\lambda_{m_d^*(\mu)}}$ so that

$$\begin{aligned} [W : W_{\mu}] &\geq [W : W_{\lambda_{m_s(\mu)}} \cap W_{\lambda_{m_d^*(\mu)}}] \\ &= \frac{(\ell + 1)!}{m_s(\mu)! m_d(\mu)! [\ell + 1 - m_s(\mu) - m_d(\mu)]!}. \end{aligned}$$

Using part (3) of Lemma 2.1, we get $[W : W_{\mu}] \leq \binom{\ell+1}{m_s(\mu) + m_d(\mu)}$. Again, Lemma 2.1 forces $m_s(\mu) + m_d(\mu) \leq \frac{n}{e}$. ■

Note that this lemma implies that $k_d = 0$ when $G = Sp(V)$ or $Spin(V)$. The last lemma in this section is a technical one:

LEMMA 2.3. *Suppose that $k_t \leq \frac{n}{e}$ and that $r = 5, 7, 11$, or 13 .*

1. *If $e \geq 3$ then $r[k_t/(r-1)] < n-2$.*
2. *If $e \geq 4$ then $r[2k_t/(r-1)] < n-2$.*

Proof. We will give a proof for the case $e \geq 3$ and $r = 7$, all other cases following from an identical argument. As $\bar{e} \geq e$, we may assume that $k_t \leq \frac{n}{3}$. By way of contradiction, assume that $r[k_t/(r-1)] \geq n-2$. As $r = 7$, $7[k_t/(r-1)] < 7(k_t/6 + 1)$. Then $n-2 < 7k_t/6 + 7 \leq 7n/18 + 7$. Thus $n < 144/11 < 14$, which is a contradiction. ■

3. WEIGHT MODULES

Let $V_{\mathbb{Z}}$ be the \mathbb{Z} -lattice generated by $\{v_i\}$ and let $\bar{V} = V_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{C}$. Moreover, assume that \bar{G} is the complex version of G so that $\bar{G} = SL(\bar{V})$, $Sp(\bar{V})$, or $Spin(\bar{V})$. Let \mathcal{L} be the adjoint module for \bar{G} , which is a complex Lie algebra, and let \mathcal{U} be the universal enveloping algebra of \mathcal{L} . We will assume

that $\{\lambda_i\}$ are the fundamental dominant weights for \mathcal{U} , that Π is the corresponding set of simple roots, and that Φ is the root system of \mathcal{U} .

Let $\{x_\alpha, \alpha \in \Phi; h_j, 1 \leq j \leq \ell\}$ be a Chevalley basis for \mathcal{L} . The Kostant \mathbb{Z} -form $\mathcal{U}_{\mathbb{Z}}$ is generated by $\{x_\alpha^m/m! \mid \alpha \in \Phi, m \in \mathbb{Z}^+\}$. Moreover, the hyperalgebra \mathcal{H} corresponding to G is defined to be $\mathcal{U}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{F}$.

Define $v_{\lambda_i} = v_1 \wedge \cdots \wedge v_i$, which is naturally an element of $\wedge^i V_{\mathbb{Z}}$ as well as of $\wedge^i V$. For convenience, we identify these two elements. Set $\bar{v}_{\lambda_i} = v_{\lambda_i} \otimes_{\mathbb{Z}} 1_{\mathbb{C}}$. Then \bar{v}_{λ_i} is a vector of weight λ_i unless $G = \text{Spin}(V)$ and $i \geq \ell - 1$.

LEMMA 3.1. $\wedge^i V$ and $\wedge^{\dim(V)-i} V^*$ contain a highest weight vector of weight λ_i unless $G = \text{Spin}(V)$ and $i \geq \ell - 1$.

Proof. By [1, Sects. 15.3, 17.3, and 19.5], $(\wedge^i \bar{V}) \cap \mathcal{H}$ is the $\mathbb{F}\mathcal{U}$ -module with highest weight λ_i , where \mathcal{H} is the intersection of kernels of contraction maps. Moreover, as $\bar{v}_{\lambda_i} \in \mathcal{H}$, it is a highest weight vector. If $\mathcal{U}_{\mathbb{Z}}$ is the Kostant \mathbb{Z} -form of \mathcal{U} , it follows that $\mathcal{U}_{\mathbb{Z}} v_{\lambda_i} \otimes_{\mathbb{Z}} \mathbb{F}$ is the Weyl module for G of highest weight λ_i . Since $v_{\lambda_i} \in \wedge^i V$, $\wedge^i V$ contains a highest weight vector of weight λ_i , and we have the first claim. The second follows from the $\mathbb{F}SL(V)$ -isomorphism $\wedge^i V \cong \wedge^{\dim(V)-i} V^*$. ■

As a direct result of this lemma we have:

PROPOSITION 3.2. M occurs as a section of

1. $V^{\otimes k_s} \otimes V^{*\otimes k_d}$ if $G = SL(V)$ or
2. $V^{\otimes k_s}$ if $G = Sp(V)$ or $\text{Spin}(V)$.

Proof. First note that if v_{μ_1} and v_{μ_2} are highest weight vectors for G of weights μ_1 and μ_2 , respectively, then $v_{\mu_1} \otimes v_{\mu_2}$ is a highest weight vector of weight $\mu_1 + \mu_2$. In particular, if \mathcal{H}^+ are the elements of \mathcal{H} corresponding to positive root elements and $x \in \mathcal{H}^+$, then $x(v_{\mu_1} \otimes v_{\mu_2}) = (xv_{\mu_1}) \otimes v_{\mu_2} + v_{\mu_1} \otimes (xv_{\mu_2}) = 0$. From the universal property of Weyl modules, it is enough to show that the modules above contain a highest weight vector of weight λ . Let $I = \{i \mid a_i \neq 0\}$ and write $M_i = \wedge^i V$ and $M'_i = \wedge^i V^*$. Consider the module $\otimes_I M_i^{a_i}$. From the above, the highest weight of this module is $\sum a_i \lambda_i = \lambda$. The claim follows by recalling the definition of k_s and k_d , and replacing $M_{\ell-i+1}$ with M'_i when $G = SL(V)$ and $i \geq \ell/2$. ■

The next lemma extends the results of Premet [6] and Suprunenko [7].

LEMMA 3.3. Let $\mu \neq 0$ be a subdominant weight of λ . If $G = Sp(V)$ and $p = 2$, assume further that $\mu = \sum b_i \lambda_i$ with $b_\ell = 0$ and that α_ℓ is not in the support of $\lambda - \mu$. Then M possesses a vector of weight μ .

Proof. By Premet's result [6], we need only consider the case where $p = 2$ and $G = Sp(V)$. Define $\Pi_0 = \{\alpha_1, \dots, \alpha_{\ell-1}\}$ and let Φ_0 be the root system generated by Π_0 . Suppose \mathcal{U}_0 is the subalgebra of \mathcal{U} corresponding

to Π_0 ; then $(\mathcal{U}_0)_{\mathbb{Z}}$ is generated by $\{x_{\alpha}^m/m! \mid \alpha \in \Phi_0, m \in \mathbb{Z}^+\}$. Note that \mathcal{U}_0 is the universal enveloping algebra of a complex Lie algebra of type $A_{\ell-1}$. Write $\Phi^+ = \{\beta_1, \dots, \beta_j\}$ such that $\beta_i \in \Phi_0^+$ if $i \leq |\Phi_0^+|$.

Now, M is a homomorphic image of $\mathcal{U}_{\mathbb{Z}} v_{\lambda} \otimes_{\mathbb{Z}} \mathbb{F}$ where v_{λ} is a highest weight vector of weight λ . Moreover, M_{μ} is the image of the space $\langle (\prod x_{-\beta_i}^{m_i}/m_i!) v_{\lambda} \mid \lambda - \mu = \sum m_i \beta_i \rangle$. From our restrictions on λ and μ , it is clear that $m_i = 0$ if $\beta_i \notin \Phi_0$. In particular, M_{μ} is a weight space contained in the image of $(\mathcal{U}_0)_{\mathbb{Z}} v_{\lambda} \otimes_{\mathbb{Z}} \mathbb{F}$. By [7], this space is non-trivial; hence M contains a vector of weight μ . ■

LEMMA 3.4. *M possesses a vector of weight $\lambda_{k_s} + \lambda_{k_d}^*$.*

Proof. Recall that $k_d = 0$ when $G = Sp(V)$ or $Spin(V)$ and that $\lambda_0 = \lambda_{\ell+1} = 0$. By the previous lemma, it is enough to show that $\lambda_{k_s} + \lambda_{k_d}^*$ is a subdominant weight of λ and that $\lambda - \lambda_{k_s} - \lambda_{k_d}^*$ is supported on $\{\alpha_1, \dots, \alpha_{\ell-1}\}$ when $G = Sp(V)$ and $p = 2$. Using the descriptions from Section 2 write $\lambda = b_i \epsilon_i$ and note that $b_1 \geq b_2 \geq \dots \geq b_m$ where $m = \max\{i \mid b_i \neq 0\}$. We will consider the case $G = Sp(V)$ or $Spin(V)$ first. Here $\sum b_i = k_s$ so that $k_s - m \geq 0$. We will induct on $k_s - m$. If $k_s - m = 0$ then $m = k_s$ and $\lambda = \lambda_{k_s}$, so the claim follows in this case. Assume $k_s - m > 0$. This implies that $b_i > 1$ for some i , so pick m_0 to be maximal subject to $b_{m_0} > 1$. By Lemma 2.2, $m < k_s < \frac{n}{e} < \ell - 1$ so that $\alpha = \epsilon_{m_0} - \epsilon_{m+1}$ is a positive root. Moreover, using the descriptions from Section 2, $\lambda - \alpha = \sum b_i \epsilon_i - \epsilon_{m_0} + \epsilon_{m+1} = \mu$ is a dominant weight. As $m+1 < \ell$, $\lambda - \mu$ is supported on $\{\alpha_1, \dots, \alpha_{\ell-1}\}$. By induction, λ_{k_s} is a subdominant weight of μ , hence of λ . Moreover, $\lambda - \lambda_{k_s}$ contains no α_{ℓ} term. Thus the result holds in this case.

Now suppose that $G = SL(V)$. Here we write $\lambda = \mu_1 + \mu_2$, where $\mu_1 = \sum_{i < \ell/2} b_i \lambda_i$ and $\mu_2 = \sum_{i \geq \ell/2} b_i \lambda_i$. The previous argument shows that λ_{k_s} is a subdominant weight of μ_1 and that λ_{k_d} is a subdominant weight of $\rho(\mu_2)$. Thus $\rho(\lambda_{k_d}) = \lambda_{k_d}^*$ is a subdominant weight of μ_2 and $\lambda_{k_s} + \lambda_{k_d}^*$ is a subdominant weight of $\mu_1 + \mu_2 = \lambda$. ■

Note that Lemma 2.2 implies that $k_s + k_d \leq \frac{n}{e}$.

4. THE SUBGROUP \tilde{C} AND SUBMODULE M_0

Assume that $r \neq p$ is an odd prime and that $k > 0$ is an integer such that $kr < n - 2$. Let H_1 be a subgroup of $H/Z(H)$ such that $H_1 \cong S_{n-2}$ and let $E_{r,k}$ be a subgroup of H_1 generated by k disjoint r -cycles. Then $E_{r,k}$ is an elementary abelian r -group of rank k . Let $C = C_{H_1}(E_{r,k})$ and $N = N_{H_1}(E_{r,k})$. Then $C = E_{r,k} C_1$ where $C_1 \cong S_{n-kr-2}$ and $N = N_1 C_1$ where $N_1 \cong \mathbb{Z}_r : \mathbb{Z}_{r-1} \wr S_k$ and $[N_1, C_1] = 1$. The action of N on $E_{r,k}$ is

easy to see; in particular, if $1 \neq \sigma \in E_{r,k}$ then $\sigma^N = \sigma^{N_1}$ and $\sigma^i \in \sigma^N$ for $(i, r) = 1$. Moreover, if σ is the product of m disjoint r -cycles, then $C_N(\sigma) \cong \mathbb{Z}_r \wr S_m \times \mathbb{Z}_r : \mathbb{Z}_{r-1} \wr S_{k-m} \times C_1$. Thus $|\sigma^N| = (r-1)^m \binom{k}{m}$. As $\binom{k}{m} \geq k$ for $0 < m < k$ and $(r-1)^{k-1} \geq k$ for all r, k as above, we have that $|\sigma^N| = (r-1)^m \binom{k}{m} \geq (r-1)k$.

Let $\tilde{E}_{r,k}$, \tilde{C} , \tilde{C}_1 , \tilde{N} , and \tilde{N}_1 be the full preimage in H of $E_{r,k}$, C , C_1 , N , and N_1 , respectively. Since $r > 2$, we may identify $E_{r,k} = O_r(\tilde{E}_{r,k})$ and note that the action of \tilde{N} on $E_{r,k}$ is the same as the action of N on $E_{r,k}$, so the above observations still hold. Define $E_{r,k}^* = \text{Hom}(E_{r,k}, \mathbb{F}^*)$. Then the \tilde{N} action on this subgroup is just the dual of its action on $E_{r,k}$. If $1 \neq \varphi \in E_{r,k}^*$, we abuse notation and write φ^i for the element $\varphi^i: \sigma \mapsto \varphi(\sigma^i)$ of $E_{r,k}^*$.

Now, $V \downarrow_{E_{r,k}} = \bigoplus_{\varphi \in E_{r,k}^*} V_\varphi$, where V_φ is the homogeneous component of φ on V . Moreover, each V_φ is an $\mathbb{F}\tilde{C}$ -module.

LEMMA 4.1. *Assume that $V_{\varphi_1}, V_{\varphi_2} \neq 0$. Then*

1. *If $\varphi_1 \in \varphi_2^{\tilde{N}}$ then $V_{\varphi_1} \cong V_{\varphi_2}$ as $\mathbb{F}\tilde{C}_1$ -modules.*
2. *If $\varphi_1 \neq 1$ then V_{φ_1} is a self-dual $\mathbb{F}\tilde{C}_1$ -module.*
3. *$V_{\varphi_1} \perp V_{\varphi_2}$ if $\varphi_1 \neq \varphi_2^{-1}$.*
4. *If $\varphi_i \neq 1$ then V_{φ_i} is totally singular.*

Proof. From the earlier comments, $\varphi_2^{\tilde{N}} = \varphi_2^{\tilde{N}_1}$ and $[\tilde{C}_1, \tilde{N}_1] = 1$. This implies that $V_{\varphi_1} \cong V_{\varphi_2}$ as $\mathbb{F}\tilde{C}_1$ -modules. Let $\sigma \in \tilde{N}_1$ be an element which interchanges φ_1 and φ_1^{-1} . As $(V_{\varphi_1})^* \cong V_{\varphi_1^{-1}}$ as an $\mathbb{F}E_{r,k}$ -module, $V_{\varphi_1} \oplus (V_{\varphi_1})^*$ is a module for $(E_{r,k} \langle \sigma \rangle) \tilde{C}_1$. As σ centralizes \tilde{C}_1 , $V_{\varphi_1} \cong V_{\varphi_1}^*$ as $\mathbb{F}\tilde{C}_1$ -modules, proving (2). We may assume that $G = Sp(V)$ or $Spin(V)$ for the remaining parts. Let \mathbf{f} be the non-degenerate, G -invariant, bilinear form on V and let Q be the associated quadratic form when $G = Spin(V)$. Suppose $u_i \in V_{\varphi_i}$ so that $gu_i = \varphi_i(g)u_i$ for all $g \in E_{r,k}$. Then $\mathbf{f}(u_1, u_2) = \mathbf{f}(gu_1, gu_2) = \varphi_1(g)\varphi_2(g)\mathbf{f}(u_1, u_2)$ for all $g \in E_{r,k}$. If $\mathbf{f}(u_1, u_2) \neq 0$ then $\varphi_1 = \varphi_2^{-1}$, hence (3) is true. (4) follows from (3) by taking $\varphi_1 = \varphi_2$, unless $G = Spin(V)$ and $p = 2$. In this case, however, $Q(u_1) = Q(gu_1) = \varphi_1^2(g)Q(u_1)$. As $r > 2$, $\varphi_1^2 = 1$ if and only if $\varphi_1 = 1$. Thus $Q(u_1) = 0$, proving (4). ■

From this point forward, we will assume that $r = 5, 7$, or 11 and $(r, p) = 1$. Also, we assume that $e \geq 3$ if $G = SL(V)$ and that $e \geq 4$ if $G = Sp(V)$ or $Spin(V)$. Let $k_t = k_s + k_d$. By the remark following Lemma 3.4, $k_t < \frac{n}{e}$. By Lemma 2.3, if $G = SL(V)$, then $r[k_t/(r-1)] < n-2$. Similarly, if $G = Sp(V)$ or $Spin(V)$, then $r[2k_t/(r-1)] < n-2$. Therefore we can take $k = [k_t/(r-1)]$ if $G = SL(V)$ and $k = [2k_t/(r-1)]$ if $G = Sp(V)$ or $Spin(V)$.

Let $1 \neq \varphi \in E_{r,k}^*$ such that $V_\varphi \neq 0$. Set $\varphi_1 = \varphi$ and write $\varphi^{\tilde{N}} = \{\varphi_1, \dots, \varphi_m\}$ such that $\varphi_i = \varphi_{m-i+1}^{-1}$, so that $\bigoplus_{i=1}^{m/2} V_{\varphi_i}$ is a totally singular subspace. Recall from the initial remarks in this section that $m \geq (r-1)k$. Hence $m \geq k_t$ when $G = SL(V)$ and $m/2 \geq k_t$ when $G = Sp(V)$ or $Spin(V)$.

By replacing T with a suitable conjugate, we may assume that each V_{φ_i} with $i \leq m/2$ has a basis which is a subset of $\{v_j\}$ where $j \leq \dim(V)/2$. Moreover, when $G = SL(V)$, we may assume that $E_{r,k} < T$. Write $V^* \downarrow_{E_{r,k}} = \bigoplus_{\psi \in E_{r,k}^*} V'_\psi$. Because $t \in T$ acts as t^{-1} on V^* and $E_{r,k} < T$, we have that $V'_\psi \cong V_{\psi^{-1}}$ as $\mathbb{F}E_{r,k}$ -modules. In particular, if $\{v_{i_1}, \dots, v_{i_j}\}$ is a basis for V_ψ then $\{v_{i_1}^*, \dots, v_{i_j}^*\}$ is a basis for V'_ψ . Moreover, $V'_\psi \cong V_\psi$ as $\mathbb{F}\tilde{C}_1$ -modules as V_ψ is a self-dual $\mathbb{F}\tilde{C}_1$ -module.

Define $U_i = V_{\varphi_i}$ if $i \leq k_s$ and $U_i = V'_{\varphi_i}$ if $k_s < i \leq k_s + k_d$.

LEMMA 4.2. $M \downarrow_{\tilde{C}}$ possesses a submodule isomorphic to $U = \bigotimes_{i=1}^{k_t} U_i$.

Proof. From Proposition 3.2, M occurs as a section of $V^{\otimes k_s} \otimes V^{*\otimes k_d}$. Thus it contains an $\mathbb{F}G$ -submodule \overline{M} such that M is a homomorphic image of \overline{M} . Let ω be the corresponding homomorphism of \overline{M} . Recall that $k_d = 0$ when $G = Sp(V)$ or $Spin(V)$.

Notice that $\bigotimes_{i \leq k_s} U_i \subset V^{\otimes k_s}$ and $\bigotimes_{i > k_s} U_i \subset V^{*\otimes k_d}$, so $U \subset V^{\otimes k_s} \otimes V^{*\otimes k_d}$. Each basic tensor in $V^{\otimes k_s} \oplus V^{*\otimes k_d}$ is a weight vector for T . Moreover, U has a basis consisting of basic tensors. Since the tensor factors in U are distinct, every basic tensor in U corresponds to a distinct weight. Pick $u_i \in U_i$ such that $u_i \in \{v_j\}$ or $u_i \in \{v_j^*\}$ and let $u = \bigotimes_{i=1}^{k_t} u_i$. It is clear that u_i is a weight vector. Because of our definition of U_i , in particular because the v_i and v_j^* that occur all have distinct indices, there is an element $w \in W$ such that $u_i = v_i$ if $i \leq k_s$ and $u_i = v_{\ell+1-i}^*$ if $k_s < i \leq k_s + k_d$. This means that u^w is a vector of weight $\lambda_{k_s} + \lambda_{k_d}^*$. Thus every element of U is conjugate, under W , to u^w .

Now recall from [3, Section 2] that the group $\mathcal{S} = \mathcal{S}_{k_s} \times \mathcal{S}_{k_d}$ acts on $V^{\otimes k_s} \otimes V^{*\otimes k_d}$ by permuting the tensor factors. A straightforward calculation, which we omit, shows that $\langle \mathbb{F}\mathcal{S}u^w \rangle$ is the entire $\lambda_{k_s} + \lambda_{k_d}^*$ weight space of $V^{\otimes k_s} \otimes V^{*\otimes k_d}$. Using Lemma 3.4, we see that $\omega(\langle \mathbb{F}\mathcal{S}u^w \rangle \cap \overline{M})$ must be a non-trivial subspace of M . Pick $\sigma \in \mathbb{F}\mathcal{S}$ such that $\sigma u^w \in \overline{M}$ and $\omega(\sigma u^w) \neq 0$. We claim that $\sigma U \subset \overline{M}$ and that $\omega|_{\sigma U}$ is injective.

First, \overline{M} is an $\mathbb{F}G$ -module containing σu^w . Thus, if w' is any element of W , then $(\sigma u^w)^{w'} = \sigma u^{ww'}$. This is because the action of G commutes with the action of \mathcal{S} . Since $u \in U$ was arbitrary and all basic vectors of U are W -conjugate to u , $\sigma U \subset \overline{M}$. Now suppose that $u' \neq u$ is any basic tensor in U . Then u' and u are vectors of distinct weights, and so are σu and $\sigma u'$. In particular, σU has a basis consisting of vectors of distinct weights.

Moreover, their images under ω remain distinct and span a subspace of M of dimension $\dim(U)$. Thus $\omega|_{\sigma U}$ is injective. Note that this implies that σU is isomorphic to U as an $\mathbb{F}\tilde{C}$ -module. As σU is an $\mathbb{F}\tilde{C}$ -module and ω is a homomorphism, we have that $\omega(\sigma U)$ is an $\mathbb{F}\tilde{C}$ -submodule of M which is isomorphic to σU , hence to U . ■

LEMMA 4.3. *Let $\delta = 0$ if k_t is even and let $\delta = 1$ if k_t is odd. Then M possesses an $\mathbb{F}\tilde{C}$ -submodule M_0 such that $\dim(M_0) \leq \ell^\delta$.*

Proof. Note that $k_t \geq 2$ so that $\dim(U_i) \leq \ell$. From Lemma 4.1, V_{φ_i} is self-dual as an $\mathbb{F}\tilde{C}_1$ -module. Similarly, V'_{φ_i} is also self-dual. Moreover, $V_{\varphi_i} \cong V'_{\varphi_i}$ as $\mathbb{F}\tilde{C}_1$ -modules. We induct on $|k_t|$. If $k_t = 1$ then the result is immediate. If $|k_t| = 2$ then $U = U_1 \otimes U_2$, where U_1 and U_2 are equivalent, self-dual $\mathbb{F}\tilde{C}_1$ -modules. Hence \tilde{C}_1 stabilizes a one-dimensional subspace \overline{M}_0 of U . If $k_t > 2$ then write $U = U_1 \otimes U_2 \otimes U'$. As U' has $k_t - 2$ tensor factors, by induction $U' \downarrow_{\tilde{C}_1}$ has a subspace \overline{M}_1 of dimension at most ℓ^δ . As $E_{r,k}$ acts as scalars on U , \tilde{C} stabilizes $\overline{M}_0 \otimes \overline{M}_1$. Take M_0 to be the image of $\overline{M}_0 \otimes \overline{M}_1$ in M under the homomorphism from Lemma 4.2. ■

5. PROOF OF THEOREM 1.1

In this section we begin the proof of Theorem 1.1. Using Lemma 4.3 and Frobenius reciprocity, we have $\dim(M) \leq \ell^\delta [H : \tilde{C}]$. In particular, $\dim(M) \leq \ell^\delta (n!/r^k(n-rk-2)!).$ And, by Lemma 3.4, $[W : W_{\lambda_{k_s} + \lambda_{k_d^*}}] \leq \dim(M)$. By combining these inequalities, we have

$$[W : W_{\lambda_{k_s} + \lambda_{k_d^*}}] \leq \ell^\delta \frac{n!}{r^k(n-rk-2)!}. \quad (1)$$

Throughout this section we will assume that $r = 7$ if $p = 5$ and $r = 5$ otherwise.

Case 1. $G = SL(V)$ and $k_t \geq 13$. Using part (3) of Lemma 2.1,

$$[W : W_{\lambda_{k_s} + \lambda_{k_d^*}}] \geq \binom{\ell+1}{k_s + k_d} = \binom{\ell+1}{k_t}.$$

Using part (1) of Lemma 2.1,

$$\binom{\ell+1}{k_t} \geq \left(\frac{\ell - k_t + 2}{k_t} \right)^{k_t}.$$

As in the previous case, $\ell - k_t + 2 < \frac{\ell}{2}$. Then using inequality (1), we get

$$\left(\frac{\ell}{2k_t} \right)^{k_t} < \ell^\delta \frac{n!}{r^k(n-rk-2)!}. \quad (2)$$

So

$$\ell^{k_t - \delta} < \frac{n!}{r^k(n - rk - 2)}(2k_t)^{k_t} < n^{rk+2}(2k_t)^{k_t}.$$

As before $\ell \geq n^3$ so that $1 < n^{rk+2-3k_t+3\delta}(2k_t)^{k_t}$. When $r = 5$,

$$rk \leq 5\frac{k_t + 3}{4} \quad \text{and} \quad rk + 2 - 3k_t + 3\delta \leq \frac{35 - 7k_t}{4}.$$

When $r = 7$,

$$rk \leq 7\frac{k_t + 5}{6} \quad \text{and} \quad rk + 2 - 3k_t + 3\delta \leq \frac{65 - 11k_t}{6}.$$

In both cases $rk + 2 - 3k_t + 3\delta < -k_t$ when $k_t \geq 14$. As $k_t < \frac{n}{e}$, $n^{rk+2-3k_t+3\delta}k_t^{k_t} < 1$ if $k_t \geq 13$, hence the result in this case.

Case 2. $G = SL(V)$ and $k_t \leq 12$. For $k_t \geq 6$, we explicitly compute k and δ for $r = 5, 7$, and 11 . We then evaluate inequality (2) for these values. In all cases, we have a contradiction for at least two primes. For $k_t = 3$ or $k_t = 4$, we assume $\ell \geq n^4$. As $8 < n$, we may use $r = 3$ as well as $r = 5$. Again using inequality (2) with these primes, we get a contradiction. For $k_t = 5, 11 < n$ so we may use $r = 3$ as well as $r = 7$. Here we use inequality (1), in particular,

$$\binom{\ell + 1}{5} > \frac{\ell^\delta}{9} \frac{n!}{(n - 11)!}.$$

Finally, the results of [4] eliminate the possibility $k_t = 2$. This completes part (1) of Theorem 1.1.

Case 3. $G = Sp(V)$ or $Spin(V)$ and $k_t \geq 16$. In this case, $k_t = k_s$ so that $[W : W_{\lambda_{k_s} + \lambda_{k_d}^*}] = 2^{k_t} \binom{\ell}{k_t}$. By part (1) of Lemma 2.1,

$$2^{k_t} \binom{\ell}{k_t} \geq 2^{k_t} \left(\frac{\ell - k_t + 1}{k_t} \right)^{k_t}.$$

Since $\ell \geq n^4$ and $k_t < \frac{n}{e}$, $\ell - k_t + 1 > \frac{\ell}{2}$. Thus

$$2^{k_t} \binom{\ell}{k_t} > \left(\frac{\ell}{k_t} \right)^{k_t}.$$

Using inequality (1), we get

$$\left(\frac{\ell}{k_t} \right)^{k_t} < \ell^\delta \frac{n!}{r^k(n - rk - 2)!}. \quad (3)$$

Then

$$\ell^{k_t-\delta} < \frac{n!}{r^k(n-rk-2)} k_t^{k_t} < n^{rk+2} k_t^{k_t}.$$

As $\ell \geq n^4$, we have $n^{4k_t-4\delta} < n^{rk+2} k_t^{k_t}$ and $1 < n^{rk+2-4k_t+4\delta} k_t^{k_t}$. When $r = 5$,

$$rk \leq 5 \frac{k_t+1}{2} \quad \text{and} \quad rk+2-4k_t+4\delta \leq \frac{13-3k_t}{2}.$$

When $r = 7$,

$$rk \leq 7 \frac{k_t+2}{3} \quad \text{and} \quad rk+2-4k_t+4\delta \leq \frac{32-5k_t}{3}.$$

In both cases $rk+2-4k_t+4\delta < -k_t$ when $k_t \geq 16$. As $k_t < \frac{n}{e}$, $n^{rk+2-4k_t+4\delta} k_t^{k_t} < 1$ if $k_t \geq 16$, hence the result in this case.

Case 4. $G = Sp(V)$ or $Spin(V)$ and $1 < k_t \leq 15$. When $k_t = 4$, 6 or $k_t \geq 8$, we explicitly compute k and δ for $r = 5$, 7, 11, and 13. We then evaluate inequality (3) for these values. In all cases we end up with a contradiction for at least two distinct values of r . We omit these calculations as they are straightforward. The remaining possibilities for k_t , namely, $k_t = 2$, 3, 5, and 7, require subtler arguments. When $k_t = 2$, it is clear that $8 < n$. Here we may take $r = 3$ or $r = 5$. These choices for r lead to a contradiction as before. When $k_t = 5$, we take $r = 5$ and this gives a contradiction, except when $p = 5$. For this case we take $r = 11$ and work with inequality (1). In particular, we have $2^5 \binom{\ell}{5} < \ell \frac{n!}{11(n-13)!}$. As $\ell \geq n^4$, this inequality leads to a contradiction. When $k_t = 7$, $n \geq 42$. Again $19, 21 < n$ so we may take $r = 17$ or $r = 19$ in this case. Here $k = 1$ and we have a contradiction. Finally, if $k_t = 3$ then $n \geq 18$. If $\ell > n^6$ then we get a contradiction using $r = 3$ and $r = 7$. This completes part (2) of Theorem 1.1.

Finally, suppose that $G = Sp(V)$, $p > 2$, and that V is an $\mathbb{F}A_n$ -module. As no irreducible $\mathbb{F}A_n$ -module has an invariant symplectic form, V must be reducible. Let V_0 be an irreducible submodule. Then, as before, there can be no invariant symplectic form on V_0 , hence it is totally singular. Thus H is contained in a parabolic subgroup of G , hence of $GL(M)$. As this is not the case, $V \downarrow_H$ must have one constituent which is a faithful $\mathbb{F}2.A_n$ -module. By [8], $\dim(V)$ is divisible by $2^{(n-s-1)/2}$ where s is the number of 1's in the binary expansion of n . In particular, $\dim(V) > 2^{(n-3)/2} / \sqrt{n}$. By parts (1) and (2) of Theorem 1.1, we have that $\dim(V) \leq 2n^6$. Comparing these bounds we see that $n < 74$. This completes the proof of Theorem 1.1. ■

REFERENCES

1. W. Fulton and J. Harris, "Representation Theory," Springer-Verlag, Berlin/New York, 1991.
2. J. E. Humphreys, "Introduction to Lie Algebras and Representation Theory," Springer-Verlag, Berlin/New York, 1972.
3. W. J. Husen, Restrictions of $\Omega_m(q)$ -modules to alternating groups, *Pacific J. Math.*, to appear.
4. K. Magaard and G. Malle, Irreducibility of alternating and symmetric squares, *Manuscripta Math.* **95** (1998), 169–180.
5. K. Magaard, G. Röhrle, and D. Testerman, On maximal subgroups of finite classical groups, preprint.
6. A. Premet, Weights of infinitesimally irreducible representations of Chevalley groups over a field of prime characteristic, *Mat. Sb.* **61** (1988), 167–183 [English translation].
7. I. Suprunenko, The invariance of the weight system of irreducible representations of algebraic groups and Lie algebras of type A_l with restricted highest weights, under reduction modulo p , *Vestsī Akad. Navuk Belarusī Ser. Fiz. Mat. Navuk* **2** (1983), 18–22 [in Russian].
8. A. Wagner, An observation on the degrees of projective representations of symmetric and alternating groups over an arbitrary field, *Arch. Math.* **29** (1977), 583–589.